

FORM PTO-1390  
OFFICE  
(REV 11/98)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK

ATTORNEY'S DOCKET NO.  
1373 WO/USTRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

U.S. APPLICATION NO (if known see 37

CFR 1.5)

10/089036

INTERNATIONAL APPLICATION NO.  
PCT/EP00/09856INTERNATIONAL FILING DATE  
05 October, 2000PRIORITY DATE CLAIMED  
05 October, 1999

## TITLE OF INVENTION

CARBON MONOXIDE SOURCE FOR THE PREPARATION OF TRANSITION METAL CARBONYL COMPLEXES

APPLICANT(S) FOR DO/EO/US

MALLINCKRODT INC.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
  2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
  3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
  4. ☒ A proper Demand for International Preliminary Examination was made by the 19<sup>th</sup> month from the earliest claimed priority date.
  5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
    - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
    - b. ☒ has been transmitted by the International Bureau.
    - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
  6. ☐ A translation of the International Application into English (35 U.S.C. 371 (c)(2)).
  7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
    - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
    - b. ☐ have been transmitted by the International Bureau
    - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
    - d. ☒ have not been made and will not be made.
  8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
  9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
  10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
- Items 11. to 16. below concern document(s) or information included:**
11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
  12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
  13. ☐ A FIRST preliminary amendment
    - ☐ A SECOND or SUBSEQUENT preliminary amendment.
  14. ☐ A substitute specification.
  15. ☒ A change of power of attorney and/or address letter.
  16. ☐ Other items or information.

10/089036

PCT/EP00/09856

1373 WO/US

- 17.
- ☒
- The following Fees are submitted:

**BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):**

Neither international preliminary examination fee (37 CFR 1.482)  
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO  
and International Search Report not prepared by the EPO or JPO.....\$1000.00  
International preliminary examination fee (37 CFR 1.482) not paid to  
USPTO but International Search Report prepared by the EPO or JPO.....\$860.00  
International preliminary examination fee (37 CFR 1.482) not paid to USPTO but  
international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$710.00  
International preliminary examination fee paid to USPTO (37 CFR 1.482)  
but all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$690.00  
International preliminary examination fee paid to USPTO (37 CFR 1.482)  
and all claims satisfied provisions of PCT Article 33 (1)-(4).....\$100.00

**ENTER APPROPRIATE BASIC FEE AMOUNT =**

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☐ 30  
months from the earliest claimed priority date (37 CFR 1.492(e)).

**CALCULATIONS** PTO USE ONLY

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total claims	26 - 20 = 6		x \$18.00	\$108.00
Independent claims	6 - 3 = 3		x \$78.00	\$234.00
MULTIPLE DEPENDENT CLAIMS(S) (if applicable)			+ \$270.00	\$ .00

**TOTAL OF ABOVE CALCULATIONS = \$ .00**

Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement  
must also be filed (Note 37 CFR 1.9, 1.27, 1.28).

**SUBTOTAL = \$1202.00**

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30  
months from the earliest claimed priority date (37 CFR 1.492(f)).

**TOTAL NATIONAL FEE = \$1202.00**

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be  
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property.

**TOTAL FEES ENCLOSED = \$1242.00**

Amount to be  
refunded: \$  
charged: \$1242.00

a. ☐ A check in the amount of \$ to cover the above fees is enclosed.

b. ☒ Please charge my Deposit Account No. 13-1160 in the amount of \$1242.00 to cover the above fees.  
A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any  
overpayment to Deposit Account No. 13-1160. A duplicate copy of this sheet is enclosed.

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

SEND ALL CORRESPONDENCE TO:

Jeffrey S. Boone  
Mallinckrodt Inc.  
675 McDonnell Boulevard  
P.O. Box 5840  
St. Louis, Missouri 63134



SIGNATURE  
Jeffrey S. Boone  
NAME  
29,284  
REGISTRATION NUMBER

10/089036

JC10 Rec'd PCT/PTO 25 MAR 2002

PATENT

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

In re Application of: Alberto, et al.	)	
International Application: PCT EP00/09856	)	
International Filing Date: 2000 October 05	)	
US Serial Number:	)	Examiner:
US Filing Date:	)	Group Art Unit:
Attorney Docket: 1373 WO/US	)	

**PRELIMINARY AMENDMENT**

This is a preliminary amendment in the US national phase filing of a PCT application.

Please cancel all pending claims and substitute the claims on the attached sheets.

The application received a favorable preliminary examination report in the PCT proceedings. The amendments are presented in order to put the claims in form for US practice.

**Authorization to Charge Deposit Account**

If it is undersigned's understanding that fees are due, a separate cover sheet setting forth the amount of the fees and an authorization to charge the fees to a deposit account will be attached. However, in the event of an error, it is requested that any fees due be charged to Deposit Account No. 13-1160.

Respectfully submitted,




Jeffrey S. Boone, J.D.  
Attorney for Applicant(s)  
Registration No. 29,284

Mallinckrodt Inc.  
675 McDonnell Boulevard  
P.O. Box 5840  
St. Louis, MO 63134  
Telephone: (314) 654-8955  
Fax: (314) 654-3156  
2002 March 19

Q:\USB\Patents\1373 US prelim amend.doc

1. Compounds of formula I



wherein  $\text{X}_1$ ,  $\text{X}_2$  and  $\text{X}_3$  are the same or different and either a Lewis base or hydride and Y is a sigma donating group.

2. Compounds as claimed in claim 1, wherein:

$\text{X}_1$  is -H;

$\text{X}_3$  and  $\text{X}_2$  are substituents which may be the same or different and are selected from the group consisting of

-H,  $-\text{NH}_x\text{R}_y$  with  $x+y=3$ , or -R, wherein R is a substituent which is bound by a carbon atom to the nitrogen or boron, respectively, and is preferably alkyl or aryl;

Y is -OH,  $-\text{OH}_2$ , -OR or -NHR, wherein R is a substituent which is bound by a carbon atom to the nitrogen or oxygen, respectively, and is preferably an alkyl or aryl, or salts thereof.

3. A compound of claim 1 wherein the compound is a borane carbonate compound in which  $\text{X}_1$ ,  $\text{X}_2$  and  $\text{X}_3$  are -H and Y is  $-\text{OH}_2$ , and/or the corresponding salts of the mono- or dideprotonated borane carbonate  $[\text{H}_3\text{BCO}_2]^{2-}$ .

4. A compound of claim 2 wherein the compound is a borane carbonate compound in which  $\text{X}_1$ ,  $\text{X}_2$  and  $\text{X}_3$  are -H and Y is  $-\text{OH}_2$ , and/or the corresponding salts of the mono- or dideprotonated borane carbonate  $[\text{H}_3\text{BCO}_2]^{2-}$ .

5. A compound of claim 1 wherein the compound is a borane amino acid (ammonia carboxy borane) in which  $\text{X}_1$  is  $\text{NH}_3$ ,  $\text{X}_2$  and  $\text{X}_3$  are -H and Y is -OH, and/or the corresponding salts of the monodeprotonated ammine borane carbonate  $[(\text{NH}_3)\text{H}_2\text{BCO}_2]^-$ .

6. A compound in claim 1 wherein the compounds are alkylated borane amino acids (trialkyl ammonia carboxy boranes) in which  $\text{X}_1$  is  $-\text{NH}_x\text{R}_y$  with  $x+y=3$ , wherein R is a substituent which is bound by a carbon atom to the nitrogen and is preferably alkyl or aryl,  $\text{X}_2$  and  $\text{X}_3$  are -H and Y is -OH.

7. A compound of claim 1 wherein  $X_1$  is an organic substituent bound by a carbon atom to boron,  $X_2$  and  $X_3$  are -H and Y is -OH<sub>2</sub>.

8. A compound of claim 1 wherein  $X_1$ ,  $X_2$  and  $X_3$  are -H and Y is OR', in which R' is a substituent bound by a carbon atom to the oxygen, in particular an alkyl, more in particular methyl or ethyl.

9. A compound of claim 1 wherein  $X_1$ ,  $X_2$  and  $X_3$  are -H and Y is NH<sub>2</sub>, NHR" or NR"<sub>2</sub>, wherein R" is a substituent bound by a carbon atom to nitrogen, in particular an alkyl, more in particular a methyl or ethyl.

10. A compound of claim 2 selected from the group consisting of the boranocarbonate derivatives [H<sub>3</sub>B-COOH<sub>2</sub>], [H<sub>3</sub>B-COOH]M, [H<sub>3</sub>B-COO]M<sub>2</sub>, Na[H<sub>3</sub>B-COOCH<sub>3</sub>], wherein M is an alkali cation; the boranocarbamates Na[H<sub>3</sub>BCONHCH<sub>3</sub>], M[H<sub>3</sub>B-CONH<sub>2</sub>], wherein M is an alkali cation; the ammine-boranocarbonates [H<sub>3</sub>N-BH<sub>2</sub>-COOH], [H<sub>3</sub>N-BH<sub>2</sub>-COO]Li, [(CH<sub>3</sub>)<sub>3</sub>N-BH<sub>2</sub>-COOH], [(CH<sub>3</sub>)<sub>2</sub>N-BH<sub>2</sub>-COOH], [(CH<sub>3</sub>)<sub>2</sub>N-BH<sub>2</sub>-COO]Li, [(CH<sub>3</sub>)<sub>2</sub>N-BH<sub>2</sub>-COOCH<sub>3</sub>] and the ammine-boranocarbamates [H<sub>3</sub>N-BH<sub>2</sub>-CONH<sub>2</sub>], [(CH<sub>3</sub>)<sub>2</sub>HN-BH<sub>2</sub>-CONHC<sub>2</sub>H<sub>5</sub>].

11. A method for preparing transition metal carbonyl complexes, wherein one or more compounds of claim 1 are used as the CO source and optionally as the reducing agent.

12. Method as claimed in claim 11, wherein the transition metal in the transition metal carbonyl complex is selected from the groups V-B to VIII-B metals.

13. Method as claimed in claim 12, wherein the transition metal in the transition metal carbonyl complex is selected from the group consisting of Vanadium (V), Chromium (Cr), Molybdenum (Mo), Tungsten (W), Manganese (Mn), Technetium (Tc), Rhenium (Re), Iron (Fe), Ruthenium (Ru), Osmium (Os), Cobalt (Co), Rhodium (Rh), Iridium (Ir) and Nickel (Ni).

14. A kit for preparing transition metal carbonyl complexes, comprising at least one compound according to claim 1 in aqueous solution, optionally one or more stabilizers, optionally one or more additional reducing agents and a buffer system.

15. A kit for preparing transition metal carbonyl complexes, comprising at least one compound according to claim 2 in aqueous solution, optionally one or more stabilizers, optionally one or more additional reducing agents and a buffer system.

16. A kit as claimed in claim 14, wherein the stabilizers are selected from the group consisting of glucoheptonate, tartrate, citrate, lactate.

17. A kit as claimed in claim 14, wherein the additional reducing agents are selected from the group consisting of boron hydrides, dithionites,  $\text{SnCl}_2$ , sulfite.

18. A kit as claimed in claim 15, wherein the additional reducing agents are selected from the group consisting of boron hydrides, dithionites,  $\text{SnCl}_2$ , sulfite.

19. A method for the preparation of borano carbonate, comprising the steps of:

- a) reacting  $\text{BH}_3 \cdot \text{THF}$  or a similar adduct in THF or a mixture of THF and another organic aprotic solvent with CO to generate  $\text{H}_3\text{BCO}$ ;
- b) passing the  $\text{H}_3\text{BCO}$  thus generated through a cold solution of a hydroxide with a mono or dikationic counter ion and an aliphatic alcohol; and
- c) after a suitable reaction time heating the alcoholic solution to precipitate the borano carbonate.

20. A method of claim 19, wherein the similar adduct is  $\text{H}_3\text{B}(\text{Et}_2\text{O})$ .

21. A method of claim 19 wherein the hydroxide is selected from the group consisting of potassium hydroxide, sodium hydroxide or tetraalkyl ammonium hydroxide.

22. A method of claim 20 wherein the hydroxide is selected from the group consisting of potassium hydroxide, sodium hydroxide or tetraalkyl ammonium hydroxide.

23. A method of claim 19 wherein the aliphatic alcohol is selected from the group consisting of methanol, ethanol and isopropanol.

24. A method of claim 20 wherein the aliphatic alcohol is selected from the group consisting of methanol, ethanol and isopropanol.

25. A method of reducing an organic substrate with  $\text{H}_3\text{BCO}$  as a reducing agent.

26. A method of claim 25 wherein the organic substrates, is selected from esters, imines or aldehydes, in water.

Rec'd PET/PTC 25 MAR 2002

CARBON MONOXIDE SOURCE FOR THE PREPARATION OF  
TRANSITION METAL CARBONYL COMPLEXES

The present invention relates to compounds that have a novel use as a carbon monoxide source and optionally as a reducing agent in the preparation of transition metal carbonyl complexes.

5 Carbonyl complexes are compounds that contain carbon monoxide as a coordinated ligand. Carbon monoxide is a common ligand in transition metal chemistry, in part due to the synergistic nature of its bonding to transition metals.

10 The bonding of CO to a metal consists of two components. The first component of the bonding is based on  $\sigma$ -donation, the overlap of a lone pair on the carbon atom with an empty d-orbital of the metal. The second component consists in  $\pi$ -back-donation from a filled d-  
15 orbital of the metal into an empty  $\pi^*$  orbital of the carbon atom of CO. This second component is called pi-backbonding or pi-backdonation.

The above described formation of carbonyl complexes with transition metals is crucial for the  
20 application of such compounds in the labeling of proteins, peptides and a variety of other compounds. For many applications these molecules are labeled by means of a so-called labeling kit which contains the necessary reagents. Current kits are based on boron hydride as the  
25 reducing agent, further contain tartrate, lactose and borate buffer, pH 11.5, and are filled with gaseous CO as the CO source. The disadvantages of these known reaction mixtures are the slow dissolution of CO into the reaction solvent resulting in a decreased yield of carbonyl  
30 complexes, the impossibility of industrial preparation of large amounts of CO filled kit vials and the slow diffusion of CO even through tightly closed vials. Moreover, the pH is rather high, which is not convenient.



It is the object of the present invention to provide an alternative for CO and sodium boron hydride that does not have the above stated drawbacks.

It has now been found that compounds of formula

5 I

10



(I)

wherein:

$X_1$  is -H;

- 15  $X_2$  and  $X_3$  are substituents which may be the same or different and are selected from the group consisting of -H,  $-NH_xR_y$  with  $x+y=3$ , or -R, wherein R is a substituent which is bound by a carbon atom to the nitrogen or boron, respectively, and is preferably alkyl or aryl;
- 20 Y is -OH,  $-OH_2$ , -OR or -NHR, wherein R is a substituent which is bound by a carbon atom to the nitrogen or oxygen, respectively, and is preferably alkyl or aryl; or salts thereof
- can be used as a carbon monoxide (CO) source and
- 25 optionally also as a reducing agent in the preparation of metal carbonyl complexes in aqueous solution. If Y is -OH or  $-OH_2$ , the compounds are acids which can be deprotonated (i.e. with NaOH). In that case, the compounds which are isolated are the salts (borano carbonate anion  $R_3B-COO^{2-}$
- 30 plus the corresponding cation, e.g.  $Li^+$ ,  $Na^+$ ,  $Ca^{2+}$ ,  $Mg^{2+}$  and others). The reducing agent function is only present if at least one of  $X_1$ ,  $X_2$  and  $X_3$  is a hydrogen. For stability reasons it is preferred that two of  $X_1$ ,  $X_2$  and  $X_3$  are -H. The carbon monoxide is released upon heating an aqueous
- 35 solution of the compound.

The advantages of the above compounds are the following. CO is produced for the first time in aqueous media under controllable conditions (pH, temperature).

Carbonyl complexes of the claimed metals can be prepared

40 in water at well defined conditions instead of organic

solvents or under high pressure and high temperature. The CO source and reducing agent can be present in the same single compound, which is convenient since reduction is practically always required for the preparation of 5 carbonyls. In case the metal to be complexed is Tc-99m or Re-188/186 kits can be produced without the demand of filling a vial with toxic and volatile CO. A major advantageous embodiment is a molecule combining the different functionalities in one compound. Such compound 10 can act as a reducing agent and as an in situ CO source, where the CO is only produced if a protic solvent (like water) or a Lewis acid is present.

By varying the substituents at the different positions various types of compounds can be obtained.

- 15 These can be subdivided in the following groups:
  1. a borane carbonate compound in which  $X_1$ ,  $X_2$  and  $X_3$  are -H and Y is  $-\text{OH}_2$ , and/or the corresponding salts of the mono- or dideprotonated borane carbonate  $[\text{H}_2\text{BCO}_2]^{2-}$ ;
  2. a borane amino acid (ammonia carboxy borane) in 20 which  $X_1$  is  $\text{NH}_3$ ,  $X_2$  and  $X_3$  are -H and Y is  $-\text{OH}$ , and/or the corresponding salts of the monodeprotonated ammine borane carbonate  $[(\text{NH}_3)\text{H}_2\text{BCO}_2]^-$ ;
  3. alkylated borane amino acids (trialkyl ammonia carboxy boranes) in which  $X_1$  is  $-\text{NH}_x\text{R}_y$  with  $x+y=3$ , wherein 25 R is a substituent which is bound by a carbon atom to the nitrogen and is preferably alkyl or aryl,  $X_2$  and  $X_3$  are -H and Y is  $-\text{OH}$ .
  4. compounds of formula I wherein  $X_1$  is an organic substituent bound by a carbon atom to boron,  $X_2$  and  $X_3$  are 30 -H and Y is  $-\text{OH}_2$ .
  5. compounds of formula I wherein  $X_1$  and  $X_2$  are organic substituents bound by a carbon atom to boron,  $X_3$  is -H and Y is  $-\text{OH}_2$ .
  6. borane carboxylic acid alkylester compounds wherein 35  $X_1$ ,  $X_2$  and  $X_3$  are as defined under 1-5 above and Y is  $\text{OR}'$ , in which R' is a substituent bound by a carbon atom to the oxygen, such as an alkyl, more in particular methyl or ethyl.

7. borane carbamate compounds wherein  $X_1$ ,  $X_2$  and  $X_3$  are as defined in 1-5 above and Y is  $NH_2$ ,  $NHR$  or  $NR^2$ , wherein  $R$  is a substituent bound by a carbon atom to nitrogen, such as an alkyl, more in particular methyl or ethyl.

Particular examples of these compounds are:  
boranocarbonate derivatives:  $[H_3B-COOH_2]$ ,  $[H_3B-COOH]M$ ,  $[H_3B-COO]M_2$ ,  $Na[H_3B-COOCH_3]$ , wherein M is an alkali cation;  
boranocarbamates:  $Na[H_3BCONHCH_3]$ ,  $M[H_3B-CONH_2]$ , wherein M is an alkali cation;  
10 ammine-boranocarbonates:  $[H_3N-BH_2-COOH]$ ,  $[H_3N-BH_2-COO]Li$ ,  $[(CH_3)_3N-BH_2-COOH]$ ,  $[(CH_3)_2HN-BH_2-COOH]$ ,  $[(CH_3)_2HN-BH_2-COO]Li$ ,  $[(CH_3)_2HN-BH_2-COOCH_3]$ ;  
ammine-boranocarbamates:  $[H_3N-BH_2-CONH_2]$ ,  $[(CH_3)_2HN-BH_2-CONHC_2H_5]$

15 The compounds of the invention can be prepared by means of or analogous to the methods as described by Burg et al., J. Am. Chem. Soc. 59, 780 (1936) for  $BH_3CO$ ; Malone et al., Inorg. Chem. 6, 817 (1967) for  $M_2[H_3B-COO]$  and  $M[H_3B-COOC_2H_5]$ ; Howe et al., Inorg. Chem. 10, 930 (1971) for  $M[H_3B-CONH_2]$ ; Spielvogel et al., J. Am. Chem. Soc. 102, 6343 (1980) for  $[H_3N-BH_2-COOH]$  and  $[(CH_3)_3N-BH_2-CONHC_2H_5]$ ; Spielvogel et al., Inorg. Chem. 23, 4322 (1984) for  $[(CH_3)_2HN-BH_2-COOCH_3]$ ; Spielvogel et al., Inorg. Chem. 23, 1776 (1984) and J. Am. Chem. Soc. 98, 5702 (1976) for  $[H_3N-BH_2-CONH_2]$ ,  $[(CH_3)_2HN-BH_2-CONHC_2H_5]$ .

The invention further relates to a method for preparing transition metal carbonyl complexes, wherein one or more of the compounds defined above are used as  
30 the CO source and optionally as the reducing agent. This method comprises in summary the release of CO from any compound of the invention, in particular from one or more of the compounds 1-7, in water or buffer due to protolysis and subsequent hydrolytic reactions.  
35 Concomitantly, the metal with which a carbonyl should be formed is reduced by the hydride substituent attached to boron. The compounds of the invention, in particular compounds 1-7, are dissolved in water or buffer and the

metal is added either as a solid or as a solution. Protonation and hydrolysis of the compounds of the invention, in particular of compounds 1-7, releases CO. At the same time, the hydrides attached to the boron (-H) 5 will reduce the metal center to a valency where the metal is able to coordinate the released CO. In that moment, carbonyl complexes are formed. The method according to the invention for preparing carbonyl complexes, thus comprises mixing the borano compounds of the invention 10 with an aqueous solution of the metal in the form of a metal-ion or (per)metallate. "Metal" as used in this application is intended to encompass all forms of the metal, i.e. also metal ions and (per)metallates.

The compounds and method of the invention are 15 suitable for the formation of any carbonyl complex, but in particular those in which the transition metal in the transition metal carbonyl complex is selected from the groups V-B to VIII-B metals. More in particular the method is suitable for preparing carbonyl complexes of 20 the following transition metals: Vanadium (V), Chromium (Cr), Molybdenum (Mo), Tungsten (W), Manganese (Mn), Technetium (Tc), Rhenium (Re), Iron (Fe), Ruthenium (Ru), Osmium (Os), Cobalt (Co), Rhodium (Rh), Iridium (Ir) and Nickel (Ni) and their radioactive isotopes.

25 Furthermore, the invention provides a kit for preparing transition metal carbonyl complexes, comprising a compound according to the invention in aqueous solution, a stabilizing agent like tartrate, glucoheptonate, lactate, citrate and a buffer system like 30 borate or phosphate. In a preferred embodiment thereof, the kit of the invention contains at least 2 mg borane carbonate, preferably in a borate buffer (pH 9.1) in an oxygen-free environment under a nitrogen atmosphere. It is preferred that the total volume of the solution after 35 addition of the radioactive metal solution does not exceed 1 ml. However, larger volumes such as 2 or 3 ml may also be useful in certain circumstances. Suitable

incubation conditions comprise heating the solution for about 20 minutes to 75°C.

The compounds of the invention can furthermore be used in water for the reduction of organic compounds with a selectivity and reactivity comparable to boron hydride or cyanoborohydride.

In addition, it was found that  $H_3BCO$  can be prepared continuously from  $H_3B \cdot THF$  and reacted in situ with an alcoholic solution of potassium hydroxide to give  $K_2[H_3BCO_2]$ . The key to the preparation is the control of the equilibrium between  $H_3BCO$  and  $H_3B \cdot THF$ : THF is selectively condensed from the gas stream at -50°C, while  $H_3BCO$  (b.p. -64°C) passes on, carried by a stream of carbon monoxide. Subsequently, this gas mixture is directly bubbled through an ethanolic solution of KOH at -78°C. Nucleophilic attack of  $[OH^-]$  at the highly electrophilic carbon in  $H_3BCO$  leads to the formation of  $K_2[H_3BCO_2]$  in high yield. If required  $H_3BCO$  itself can be isolated in a cold trap at -78°C. This method of preparing  $H_3BCO$  is simpler and more convenient than the high pressure or ether-catalyzed procedures and can be scaled up to quantities of several grams or larger.

Thus, the invention relates to method for the preparation of borano carbonate, comprising the steps of:

- a) reacting  $BH_3 \cdot THF$  or a similar adduct in THF or a mixture of THF and another organic aprotic solvent with CO to generate  $H_3BCO$ ;
- b) passing the  $H_3BCO$  thus generated through a cold solution of a hydroxide with a mono or dikationic counter ion and an aliphatic alcohol; and
- c) after a suitable reaction time heating the alcoholic solution to precipitate the borano carbonate. The similar adduct is for example  $H_3B(Et_2O)$ . The hydroxide is for example selected from the group consisting of potassium hydroxide, sodium hydroxide or tetraalkyl ammonium hydroxide. The aliphatic alcohol can be selected from the group consisting of methanol, ethanol and isopropanol.

The compound  $H_3BCO$  is also part of this invention. It has reducing properties and can be used for that purpose for example in the preparation of carbonyl complexes without high pressure CO as described above but then in aprotic or only weakly protic solvents. It is also possible to use  $H_3BCO$  in situ while it is produced when CO is bubbled through THF solutions of "metals", such as for the synthesis of the macroscopic  $[TcCl_3(CO)_3]_2$  or Re analogue.

The use of the compounds according to the invention is more broadly applicable than solely for the preparation of carbonyl complexes, but can also be applied in other circumstances wherein a CO source in aqueous solution is required. The invention also relates to the use of borano carbonate or derivatives thereof as a reducing agent for organic substrates, such as esters, imines or aldehydes, in water. The reducing power of these compounds is comparable to  $BH_4^-$  or cyanoborohydride and they can thus be a substitute for e.g. cyanoborohydride in bulk industrial processes.

The present invention is further illustrated in the following examples, that are given for illustration purposes only.

25

## EXAMPLES

### EXAMPLE 1

#### Preparation of $K_2H_3BCO_2$

##### 1. Synthesis of $BH_3 \cdot CO$

4 g of  $NaBH_4$  was carefully added to 15 ml of concentrated  $H_3PO_4$  (dried overnight under high vacuum at room temperature) in vacuo (1 mbar) under vigorous stirring over a period of 2 hours. The evolving  $BH_3$  was dried by passing it through a cool trap at  $-78^\circ C$  and was condensed in a second cool trap at  $-200^\circ C$  containing 70 ml of dry DME. The second trap was disconnected from the first trap and the vacuum line. The temperature was brought to  $-40^\circ C$ . Subsequently the trap was pressurized

with 1.3 bar of dry CO. The reaction mixture was stirred in a cool bath at  $-40^{\circ}\text{C}$  (dry ice with acetonitrile) under 1.3 bar of CO overnight.

5 2. Synthesis of  $\text{K}_2\text{H}_3\text{BCO}_3$

The gas outlet of the trap was connected to a 100 ml two-neck round-bottom flask (equipped with gas inlet and reflux condenser) containing 50 ml of dry ethanol and 3 g KOH. The cool bath of the trap was removed and the evolving  $\text{BH}_3\cdot\text{CO}$  was bubbled slowly through the ethanolic KOH solution at  $0^{\circ}\text{C}$ . The DME solution was slowly heated to  $80^{\circ}\text{C}$  and the trap subsequently three times flushed with CO. After the evolution of  $\text{BH}_3\cdot\text{CO}$  had stopped the ethanolic solution was refluxed for 30 min. After cooling the solution to room temperature  $\text{K}_2\text{H}_3\text{BCO}_3$  precipitated as a white powder which was filtered by a sintered glass filter, washed with ice cold ethanol and dried under vacuum.

20

**EXAMPLE 2**

Labeling experiment using a lyophilized kit

A labeling kit was prepared by lyophilizing 1 mg  $\text{K}_2[\text{BH}_3\text{COO}]$  in 0.1 ml of 0.1M PBS, pH 7.5 in a vial that was flushed with  $\text{N}_2$ . As an alternative a 0.1M borate buffer, pH 8.5 can be used.

For labeling, 1 ml of a generator eluted  $[\text{}^{99\text{m}}\text{TcO}_4]^-$  saline solution is added. It was found that the yield is independent of the absolute amount of  $[\text{}^{99\text{m}}\text{TcO}_4]^-$ . The solution thus obtained is heated to  $75^{\circ}\text{C}$  for 20 min.

The yields are between 80 and 100% (trace 1 in Fig. 1) for pH 7.5; trace 3 in Fig. 1 for pH 8.5.

To establish the identity of the compound, picolinic acid was added directly to the reaction solution, in which the carbonyl complex had been prepared. HPLC revealed the complex  $[\text{}^{99\text{m}}\text{Tc}(\text{OH}_2)(\text{pic})(\text{CO})_3]$  (Fig. 1, trace 2) by comparison with "cold" material, in the present case the same complex made with "cold"

Rhenium. The "hot" material is found by means of a radioactivity detector, whereas the "cold" material is detected with a UV detector.

5

**EXAMPLE 3**Labeling experiment with a so-called "wet kit"

A vial containing 2 mg borane carbonate and a generator eluate of pertechnetate in borate buffer, pH 9.1, in a total volume of 1 ml was heated for 20 min. to 75°C. The labeling yield of the product [ $^{99m}\text{Tc}(\text{O}_2)(\text{CO})_3$ ] $^+$  thus obtained was higher than 97%.

15 **EXAMPLE 4**Preparation of potassium hydrogen (carboxylato)-trihydroborate starting from H.B.THF

The apparatus used consisted of a 250 ml three-necked round-bottomed flask, connected to a cold-trap by a glass tube. The other two necks of the flask were sealed with rubber septa. A PTFE tube for the introduction of gas passed into the flask. From the outlet of the cold-trap, a PTFE tube passed into a 400 ml Schlenk tube. From the side-arm of the Schlenk tube, passed a polytene tube leading to a silicone oil bubbler, which isolated the apparatus from the atmosphere.

The cold-trap and the Schlenk tube were immersed in Dewar flasks containing isopropanol. The apparatus was flushed with dry oxygen-free nitrogen for 30 30 minutes while the cold trap was cooled to -50°C and the Schlenk tube to -78°C by addition of dry ice to the respective Dewar flasks.

A solution of 5.0 g potassium hydroxide in 200 ml absolute ethanol was added to the Schlenk tube and allowed to cool to -78°C. The apparatus was briefly flushed with carbon monoxide, and 30 ml of a 1 moldm<sup>-3</sup> solution of borane-tetrahydrofuran complex in tetrahydrofuran was introduced into the round-bottomed



flask. Carbon monoxide was bubbled into the solution so that approximately one bubble per second left the apparatus via the oil bubbler. The temperature of the middle cold-trap was maintained at between -45°C and -55°C by occasional addition of dry ice.

After two hours passage of carbon monoxide, 20 ml dimethoxyethane was introduced into the round-bottomed flask and an additional 20 ml dimethoxyethane was introduced into the middle cold-trap. Carbon monoxide was passed through the apparatus as before. After one hour, the Schlenk tube was disconnected from the rest of the apparatus, and allowed to warm to room temperature. The alcoholic solution was heated under reflux for 45 minutes. The resulting white precipitate was filtered off, washed with two 5 ml portions of absolute ethanol, and dried in vacuo to give 1.26 g product (43% based on  $\text{BH}_3 \cdot \text{THF}$ ) as a white powder. Found K, 38.85% (gravimetric as  $\text{K}_2\text{Na}[\text{Co}(\text{NO}_2)_6]$ );  $\text{CH}_3\text{BKO}_2$  requires K, 39.9%.  $\delta_{\text{H}}$  (200 MHz,  $\text{D}_2\text{O}$ , 25°C) 0.80 (1:1:1:1 quartet,  $^1\text{J}(\text{H}-^{13}\text{B}) = 80 \text{ Hz}$ ; 1:1:1:1:1:1:1 septet,  $^1\text{J}(\text{H}-^{10}\text{B}) = 27 \text{ Hz}$ ).


#### EXAMPLE 5

##### Reduction of the organic substrate sodium benzaldehyde-2-sulfonate with boranocarbonate in water

Potassium boranocarbonate (100 mg) and sodium benzaldehyde-2-sulfonate (40 mg) were mixed in water (1 ml) and left to stand for 30 min at room temperature. Quantitative formation of sodium 2-(hydroxymethyl)-benzene sulfonate was confirmed by the disappearance of the  $^1\text{H}$ -NMR signal of the starting material at  $\delta = 10.77$ , and the appearance of the product signal at  $\delta = 5.04$ . The reaction mixture was odorless at the end of the experiment, indicating that the sulfonate group had not been reduced.

## CLAIMS

## 1. Compounds of formula I



wherein  $X_1$ ,  $X_2$  and  $X_3$  are the same or different and either  
 10 a Lewis base or hydride and Y is a sigma donating group.

## 2. Compounds as claimed in claim 1, wherein:

$X_1$  is -H;

$X_2$  and  $X_3$  are substituents which may be the same or  
 different and are selected from the group consisting of

15 -H,  $-NH_xR_y$  with  $x+y=3$ , or -R, wherein R is a substituent  
 which is bound by a carbon atom to the nitrogen or boron,  
 respectively, and is preferably alkyl or aryl;

Y is -OH,  $-OH_2$ , -OR or -NHR, wherein R is a substituent  
 which is bound by a carbon atom to the nitrogen or

20 oxygen, respectively, and is preferably an alkyl or aryl,  
 or salts thereof

for use as a carbon monoxide (CO) source and optionally  
 as a reducing agent in the preparation of metal carbonyl  
 complexes in aqueous solution.

25 3. Compound as claimed in claim 1 or 2 wherein  
 the compound is a borane carbonate compound in which  $X_1$ ,  
 $X_2$  and  $X_3$  are -H and Y is  $-OH_2$ , and/or the corresponding  
 salts of the mono- or dideprotonated borane carbonate  
 $[H_3BCO_2]^{2-}$ .

30 4. Compound as claimed in claim 1 or 2 wherein  
 the compound is a borane amino acid (ammonia carboxy  
 borane) in which  $X_1$  is  $NH_3$ ,  $X_2$  and  $X_3$  are -H and Y is -OH,  
 and/or the corresponding salts of the monodeprotonated  
 ammine borane carbonate  $[(NH_3)H_2BCO_2]^-$ .

35 5. Compounds as claimed in claim 1 or 2 wherein  
 the compounds are alkylated borane amino acids (trialkyl  
 ammonia carboxy boranes) in which  $X_1$  is  $-NH_xR_y$  with  $x+y=3$ ,  
 wherein R is a substituent which is bound by a carbon

atom to the nitrogen and is preferably alkyl or aryl,  $X_2$  and  $X_3$  are -H and Y is -OH.

6. Compound as claimed in claim 1 or 2 wherein  $X_1$  is an organic substituent bound by a carbon atom to boron,  $X_2$  and  $X_3$  are -H and Y is -OH<sub>2</sub>.

7. Compounds as claimed in claim 1 or 2 wherein  $X_1$ ,  $X_2$  and  $X_3$  are as defined in claims 2-6 and Y is OR', in which R' is a substituent bound by a carbon atom to the oxygen, in particular an alkyl, more in particular methyl or ethyl.

8. Compounds as claimed in claim 1 or 2 wherein  $X_1$ ,  $X_2$  and  $X_3$  are as defined in claims 2-6 and Y is NH<sub>2</sub>, NHR" or NR" <sub>2</sub>, wherein R" is a substituent bound by a carbon atom to nitrogen, in particular an alkyl, more in particular a methyl or ethyl.

9. Compounds as claimed in claim 1 or 2, selected from the group consisting of the boranocarbonate derivatives [H<sub>3</sub>B-COOH<sub>2</sub>], [H<sub>3</sub>B-COOH]M, [H<sub>3</sub>B-COO]M<sub>2</sub>, Na[H<sub>3</sub>B-COOCH<sub>3</sub>], wherein M is an alkali cation; the boranocarbamates Na[H<sub>3</sub>BCONHCH<sub>3</sub>], M[H<sub>3</sub>B-CONH<sub>2</sub>], wherein M is an alkali cation; the ammine-boranocarbonates [H<sub>3</sub>N-BH<sub>2</sub>-COOH], [H<sub>3</sub>N-BH<sub>2</sub>-COO]Li, [(CH<sub>3</sub>)<sub>3</sub>N-BH<sub>2</sub>-COOH], [(CH<sub>3</sub>)<sub>2</sub>N-BH<sub>2</sub>-COOH], [(CH<sub>3</sub>)<sub>2</sub>N-BH<sub>2</sub>-COO]Li, [(CH<sub>3</sub>)<sub>2</sub>N-BH<sub>2</sub>-COOCH<sub>3</sub>] and the ammine-boranocarbamates [H<sub>3</sub>N-BH<sub>2</sub>-CONH<sub>2</sub>], [(CH<sub>3</sub>)<sub>2</sub>NH-BH<sub>2</sub>-CONHC<sub>2</sub>H<sub>5</sub>].

10. Method for preparing transition metal carbonyl complexes, wherein one or more compounds as claimed in claims 1-13 are used as the CO source and optionally as the reducing agent.

11. Method as claimed in claim 10, wherein the transition metal in the transition metal carbonyl complex is selected from the groups V-B to VIII-B metals.

12. Method as claimed in claim 11, wherein the transition metal in the transition metal carbonyl complex is selected from the group consisting of Vanadium (V), Chromium (Cr), Molybdenum (Mo), Tungsten (W), Manganese (Mn), Technetium (Tc), Rhenium (Re), Iron (Fe), Ruthenium (Ru), Osmium (Os), Cobalt (Co), Rhodium (Rh), Iridium (Ir) and Nickel (Ni).

13. Kit for preparing transition metal carbonyl complexes, comprising at least one compound according to claims 1-9 in aqueous solution, optionally one or more stabilizers, optionally one or more additional reducing agents and a buffer system.

14. Kit as claimed in claim 13, wherein the stabilizers are selected from the group consisting of glucoheptonate, tartrate, citrate, lactate.

15. Kit as claimed in claim 13 or 14, wherein the additional reducing agents are selected from the group consisting of boron hydrides, dithionites,  $\text{SnCl}_2$ , sulfite.

16. Method for the preparation of borano carbonate, comprising the steps of:

17. a) reacting  $\text{BH}_3 \cdot \text{THF}$  or a similar adduct in THF or a mixture of THF and another organic aprotic solvent with CO to generate  $\text{H}_3\text{BCO}$ ;

b) passing the  $\text{H}_3\text{BCO}$  thus generated through a cold solution of a hydroxide with a mono or dikationic counter ion and an aliphatic alcohol; and

c) after a suitable reaction time heating the alcoholic solution to precipitate the borano carbonate.

18. Method as claimed in claim 16, wherein the similar adduct is  $\text{H}_3\text{B}(\text{Et}_2\text{O})$ .

19. Method as claimed in claim 16 or 17, wherein the hydroxide is selected from the group consisting of potassium hydroxide, sodium hydroxide or tetraalkyl ammonium hydroxide.

20. Method as claimed in claim 16 or 17, wherein the aliphatic alcohol is selected from the group consisting of methanol, ethanol and isopropanol.

21. Method as claimed in claims 16-19, wherein the method is performed as described in example 4.

22. Use of  $\text{H}_3\text{BCO}$  as a reducing agent.

23. Use of borano carbonate or derivatives thereof as a reducing agent for organic substrates, such as esters, imines or aldehydes, in water.

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number  
WO 01/25243 A1

(51) International Patent Classification<sup>7</sup>: C07F 5/02, 19/00

(21) International Application Number: PCT/EP00/09856

(22) International Filing Date: 5 October 2000 (05.10.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
99203254.0 5 October 1999 (05.10.1999) EP

(71) Applicant (for all designated States except US):  
MALLINCKRODT INC. [US/US]; 675 McDonell  
Boulevard, St. Louis, MO 63134 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): ALBERTO, Roger,  
Ariel [CH/CH]; St. Georgenstrasse 54, CH-8400 Win-  
terthur (CH).

(74) Agent: VAN SOMEREN, Petronella, Francisca, Hen-  
drika, Maria; Arnold & Siedsma, Sweelinckplein 1,  
NL-2517 GK The Hague (NL).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CARBON MONOXIDE SOURCE FOR PREPARATION OF TRANSITION-METAL-CARBONYL-COMPLEXES



(57) Abstract: The present invention relates to compounds that have a novel use as a carbon monoxide source and optionally as a reducing agent in the preparation of transition metal carbonyl complexes. The compounds are in general compounds of formula (I) wherein  $X_1$ ,  $X_2$  and  $X_3$  are the same or different and either a Lewis base or hydride and Y is a sigma donating group. The invention furthermore relates to a method for the preparation of borane carbonate and to the use of  $H_3BCO$  as a reducing agent.

WO 01/25243 A1

## DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

As a below named inventor, I declare that:

My residence and post office address and my citizenship are as stated below next to my name.

I believe I am the original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled: **"CARBON MONOXIDE SOURCE FOR THE PREPARATION OF TRANSITION METAL CARBONYL COMPLEXES"** the Specification of which was filed on October 5, 2000 as PCT International Application Serial Number PCT/EP00/09856.

I hereby state that I have reviewed and understand the contents of the above identified Specification, including the Claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR 1.56, including for continuation-in-part applications, material information which became available between the filing date of the prior application and the national or PCT international filing date of the continuation-in-part application.

I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or (f), or 365(b) of any foreign application(s) for patent, inventor's or plant breeder's rights certificate(s), or 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent, inventor's or plant breeder's rights certificate(s), or any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Foreign Filing Date (MM/DD/YYYY)	Priority Not Claimed	Certified Copy Attached?	
				YES	NO
99203254.0	Europe	10/05/1999	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
			<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

I hereby appoint the following attorneys to prosecute this application and to transact all business in the Patent & Trademark Office connected therewith.

Jeffrey S. Boone  
Lawrence L. Limpus

Reg. No. 29,284  
Reg. No. 27,816

(2)

Further, I hereby request that all correspondence about the present application be addressed to:

Jeffrey S. Boone  
~~Mallinckrodt Inc.~~  
675 McDonnell Blvd.  
P.O. Box 5840  
St. Louis, Missouri 63134

I hereby certify that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Full Name of Joint Inventor:

Roger ALBERTO

Signature of Inventor

Residence:

St. Georgenstrasse 54  
Winterthur, Switzerland  
CH-8400

Post Office Address:

Same as Residence

Full Name of Joint Inventor:

Roger SCHIBLI

Signature of Inventor

Residence:

Weite Gasse 28  
Baden, Switzerland  
CH-5400

5.02.02

Date

Switzerland

Citizenship

11.02.02

Date

Switzerland

Citizenship